

Twisted-mass lattice QCD with mass non-degenerate quarks *

R. Frezzotti^a and G.C. Rossi^b

^aINFN, Sezione di Milano and Dipartimento di Fisica, Università di Milano “*Bicocca*”
Piazza della Scienza 3 - 20126 Milano (Italy)

^bDipartimento di Fisica, Università di Roma “*Tor Vergata*” and INFN, Sezione di Roma 2
Via della Ricerca Scientifica - 00133 Roma (Italy)

The maximally twisted lattice QCD action of an $SU_f(2)$ doublet of mass degenerate Wilson quarks gives rise to a real positive fermion determinant and it is invariant under the product of standard parity times the change of sign of the coefficient of the Wilson term. The existence of this spurionic symmetry implies that $O(a)$ improvement is either automatic or achieved through simple linear combinations of quantities taken with opposite external three-momenta. We show that in the case of maximal twist all these nice results can be extended to the more interesting case of a mass non-degenerate quark pair.

1. Introduction

It has been shown in ref. [1] that in lattice QCD (LQCD) with mass degenerate $SU_f(2)$ doublets of Wilson fermions it is possible to improve the approach to the continuum limit of correlation functions of gauge invariant multiplicative renormalizable (m.r.) operators by taking the arithmetic average (Wilson average - *WA*) of pairs of correlators computed in theories regularized with Wilson terms of opposite sign and identical values of the subtracted (unrenormalized) lattice quark mass, $m_q^W = M_0 - M_{cr}$ with M_0 the bare quark mass. Equivalently one can take appropriate linear combinations of the correlators computed in a given regularization (fixed sign of the Wilson term) but with opposite values of m_q^W (mass average - *MA*). Improved hadronic masses and matrix elements can be similarly obtained by taking *WA*'s of the corresponding quantities separately computed within the two regularizations.

To avoid the difficulties related to the nature of the spectrum of the Wilson-Dirac (WD) operator (here we are referring to the well-known problem of the existence of “exceptional configurations” [2,3]) twisted-mass lattice QCD (tm-LQCD) [4] should be used for the actual com-

putation of the correlators taking part to the averages. The fermionic determinant of tm-LQCD is, in fact, real and positive on arbitrary gauge backgrounds, as long as the quark mass is non-vanishing. All the nice cancellations of $O(a)$ terms that one finds in the standard Wilson case extend to tm-LQCD with mass degenerate quark doublets.

Peculiar simplifications occur if the twisting angle is taken to be equal to $\pi/2$ (maximal twist). This choice is particularly useful for applications because essentially all interesting physical quantities (e.g. hadronic masses and matrix elements of m.r. operators between states with vanishing three-momenta) can be extracted from $O(a)$ improved lattice data without making recourse to any *WA*.

The main goal of this note is to show that in the maximally twisted case positivity of the determinant and $O(a)$ improvement without *WA* can be extended to encompass the more interesting situation in which mass non-degenerate quark pairs are considered. This is a preliminary, necessary step if one wants to set up a realistic computational scheme for operator matrix elements.

For future applications it is important to stress that the approach we propose is very flexible as it allows to regularize different flavours with dif-

*Talk presented by G.C. Rossi.

ferently twisted Wilson terms. As discussed in detail in [5], it is possible to exploit this freedom in order to improve the chiral behaviour of lattice correlators to the point of (hopefully) killing all the unwanted “wrong chirality mixings” that affect in the standard Wilson [6] and twisted-mass [4,7] regularizations of QCD the construction of m.r. operators. Of special importance is the application of this new strategy to the construction of the m.r. operators which represent the CP-conserving, $\Delta S = 2$ and $\Delta S = 1$ effective weak Hamiltonian on the lattice.

The presentation of the material is divided in two parts. In the first part (sect. 2) we spell out in detail the formulae relevant for mass degenerate quark pairs with the choice $\pm\pi/2$ (maximal twist) for the twisting angle. In the second part (sect. 3) we discuss how one can deal with mass non-degenerate pairs of quarks without losing $O(a)$ improvement, or the positivity of the fermion determinant. The latter property, which is obvious in the case of mass degenerate doublets, is crucial for being able to actually carry out numerical Monte Carlo simulations. Conclusions can be found in sect. 4. In Appendix A we prove the renormalizability of the maximally twisted fermion action both in the case of mass degenerate and mass non-degenerate quarks. The proof of the positivity of the fermion determinant in the non-trivial case of mass non-degenerate quarks is given in Appendix B.

2. A summary of tm-LQCD @ $\pm\pi/2$

The twisted-mass lattice action of an $SU_f(2)$ flavour doublet of mass degenerate quark has the form

$$S_{F,D}^{(\omega)}[\psi, \bar{\psi}, U] = a^4 \sum_x \bar{\psi}(x) \left[\gamma \cdot \tilde{\nabla} + \right. \quad (1) \\ \left. + e^{-i\omega\gamma_5\tau_3} W_{\text{cr}}(r) + m_q \right] \psi(x),$$

where

$$\gamma \cdot \tilde{\nabla} \equiv \frac{1}{2} \sum_{\mu} \gamma_{\mu} (\nabla_{\mu}^* + \nabla_{\mu}), \quad (2)$$

$$W_{\text{cr}}(r) \equiv -a \frac{r}{2} \sum_{\mu} \nabla_{\mu}^* \nabla_{\mu} + M_{\text{cr}}(r) \quad (3)$$

and $M_{\text{cr}}(r) = -M_{\text{cr}}(-r)$ is the critical quark mass. We wrote the action in what is usually called the “physical basis” [1], where m_q is taken to be real (and positive)². By undoing the twisting of the Wilson term and bringing it fully to the mass term, it was proved in ref. [1] that the parameter $M_{\text{cr}}(r)$ appearing in the action (1) is equal to that of the corresponding standard Wilson theory.

In this note we will concentrate on the maximal twist case $\omega = \pi/2$, where eq. (1) becomes³

$$S_{F,D}^{(\pi/2)}[\psi, \bar{\psi}, U] = a^4 \sum_x \bar{\psi}(x) \left[\gamma \cdot \tilde{\nabla} + \right. \quad (4) \\ \left. -i\gamma_5\tau_3 W_{\text{cr}}(r) + m_q \right] \psi(x).$$

The choices $\omega = \pm\pi/2$ are particularly useful because it can be proved [1] that, despite the fact that the theory is not fully $O(a)$ improved, cancellation of $O(a)$ effects in quantities of physical interest (like energies and operator matrix elements) is either automatic or obtainable with no need of any WA . The proof of this statement is sketched in sect. 2.2. $O(a)$ ambiguities in the knowledge of $M_{\text{cr}}(r)$ do not spoil any of the above results.

Furthermore it is important to observe that at $\omega = \pm\pi/2$ the critical WD operator (i.e. the operator in square parenthesis in eq. (4) with $m_q = 0$) is anti-Hermitian, so its spectrum is purely imaginary. This means that the full WD operator cannot have any vanishing eigenvalue as soon as $m_q \neq 0$. This is also evident from the explicit expression of its determinant, $\mathcal{D}_{F,D}$, which takes the remarkably simple form

$$\mathcal{D}_{F,D} = \det \left[Q_{\text{cr}}^{\dagger} Q_{\text{cr}} + m_q^2 \right], \quad (5)$$

where $Q_{\text{cr}} = Q_{\text{cr}}^{\dagger}$ is defined by eqs. (2) and (69).

In Appendix A we prove that the fermionic action (4) is “stable” under radiative corrections, in the sense that the unbroken (possibly spurious) symmetries of the action prevent radiative corrections from generating extra independent operators (of dimension ≤ 4), not already

²Unless differently stated, we employ here the notations of ref. [1].

³Whatever we say for $\omega = \pi/2$ also holds for $\omega = -\pi/2$.

present in (4), which one would need to include for renormalizability. Once $M_{\text{cr}}(r)$ has been set to the appropriate value, the continuum limit is approached as usual by rescaling g_0^2 and m_q according to the chosen renormalization conditions.

2.1. Non-singlet Ward-Takahashi identities

In this section we collect the expressions of the renormalized current and quark density operators entering the flavour non-singlet Ward-Takahashi identities (WTI's) associated with the action (4).

Renormalized vector and axial currents can be taken to be

$$\begin{aligned}\hat{V}_\mu^1 &= Z_A \bar{\psi} \gamma_\mu \frac{\tau_1}{2} \psi & \hat{A}_\mu^1 &= Z_V \bar{\psi} \gamma_\mu \gamma_5 \frac{\tau_1}{2} \psi \\ \hat{V}_\mu^2 &= Z_A \bar{\psi} \gamma_\mu \frac{\tau_2}{2} \psi & \hat{A}_\mu^2 &= Z_V \bar{\psi} \gamma_\mu \gamma_5 \frac{\tau_2}{2} \psi \\ \hat{V}_\mu^3 &= Z_V \bar{\psi} \gamma_\mu \frac{\tau_3}{2} \psi & \hat{A}_\mu^3 &= Z_A \bar{\psi} \gamma_\mu \gamma_5 \frac{\tau_3}{2} \psi,\end{aligned}\quad (6)$$

where to make contact with known quantities the (finite) renormalization constants, Z_V and Z_A , introduced above are those for the local vector and axial currents of standard Wilson fermions, respectively. Notice the switch between Z_V and Z_A for the currents with flavour $b = 1, 2$, due to the presence of the factor $\gamma_5 \tau_3$ in front of the Wilson term in eq. (4) (see also the comment at the end of this section).

With reference to eqs. (6), the non-singlet WTI's with the insertion of the renormalized (multi-local) operator $\hat{O}(y)$ ($y \equiv \{y_i, i = 1, \dots, n\} \neq x$) take the expected form ($b = 1, 2, 3$)

$$\langle \partial_\mu^* \hat{V}_\mu^b(x) \hat{O}(y) \rangle \Big|_{(r, m_q)} = \text{O}(a) \quad (7)$$

$$\langle [\partial_\mu^* \hat{A}_\mu^b(x) - 2\hat{m}_q \hat{P}^b(x)] \hat{O}(y) \rangle \Big|_{(r, m_q)} = \text{O}(a) \quad (8)$$

with

$$\hat{m}_q = Z_P^{-1} m_q, \quad (9)$$

provided one defines, in terms of bare quantities, the renormalized pseudo-scalar operators, $\hat{P}^b$, to be

$$\hat{P}^b = Z_P \bar{\psi} \frac{\tau_b}{2} \gamma_5 \psi \quad b = 1, 2 \quad (10)$$

$$\hat{P}^3 = Z_{S^0} \left[\bar{\psi} \frac{\tau_3}{2} \gamma_5 \psi + a^{-3} i \rho_P(am_q) \mathbb{1} \right]. \quad (11)$$

In eq. (11) the dimensionless real coefficient, $\rho_P(am_q)$, in front of the power divergent term

admits a polynomial expansion in am_q . Similarly to Z_V and Z_A , Z_P and Z_{S^0} are the (logarithmically divergent) renormalization constants of the non-singlet pseudo-scalar and singlet scalar quark density of the standard Wilson regularization, respectively.

For completeness we also give the expression of the renormalized singlet scalar quark density operator, S^0 . By standard symmetry and dimensionality arguments, one finds

$$\hat{S}^0 = Z_P \left[\bar{\psi} \psi + a^{-2} m_q \rho_{S^0}(am_q) \mathbb{1} \right], \quad (12)$$

where ρ_{S^0} is a dimensionless real coefficient with an even polynomial dependence on am_q . This parity property follows from the invariance of the action (4) under the spurionic symmetry $\mathcal{P}_{\pi/2}^1 \times (m_q \rightarrow -m_q)$ (eq. (62)). Formula (12) is rather interesting because it shows that the chiral order parameter is only affected by an m_q/a^2 power divergence, analogously to what happens with Ginsparg-Wilson fermions.

The equations of this section can be obtained by specializing to the case $\omega_r = \omega = \pi/2$ the corresponding formulae of ref. [1]. A direct and practical method to deduce them is to proceed in the following way. One starts by rotating the fermionic fields in the action (4) so as to have the Wilson term in the standard form (i.e. with no factor $\gamma_5 \tau_3$ in front of it). At this point one can straightforwardly adapt to this instance the analysis of the structure of the mixing pattern of the chiral rotation of the Wilson term spelled out in detail in ref. [6] (see also ref. [8]). In a mass independent renormalization scheme renormalization constants can be computed in the massless limit ($m_q = 0$). For this reason it seemed to us a natural choice to keep for them the names they would have in the standard Wilson regularization. The final step of this method requires to rotate all the quark fields back into the physical basis we started from.

2.2. $\text{O}(a)$ improvement

It was proved in ref. [1] that the invariance of the action (1) under

$$\mathcal{R}_5^{\text{sp}} = \mathcal{R}_5 \times (r \rightarrow -r) \times (m_q \rightarrow -m_q), \quad (13)$$

$$\mathcal{R}_5 \times \mathcal{D}_d, \quad (14)$$

with $\mathcal{R}_5$ and $\mathcal{D}_d$ defined in eqs. (64) and (65), respectively, implies the validity of the formula

$$\begin{aligned} \langle O \rangle \Big|_{(m_q)}^{(\omega)WA} &\equiv \frac{1}{2} \left[\langle O \rangle \Big|_{(r, m_q)}^{(\omega)} + \langle O \rangle \Big|_{(-r, m_q)}^{(\omega)} \right] = \\ &= \zeta_O^O(\omega, r) \langle O \rangle \Big|_{(m_q)}^{\text{cont}} + \mathcal{O}(a^2), \end{aligned} \quad (15)$$

where O is any gauge invariant m.r. (multi-local) operator.

From (15) one can prove a number of physically interesting improvement formulae. To see this let us introduce the eigenstates $|h, n, \mathbf{k}\rangle_{(r, m_q)}^{(\omega)}$ (h and $\mathbf{k}$ represent the set of quantum numbers and three-momenta characterizing the state and n the excitation level) of the transfer matrix, $\hat{T}(\omega, r, m_q)$, with eigenvalues $E_{h, n}(\mathbf{k}; \omega, r, m_q)$ ⁴. The eigenvalue equation reads

$$\begin{aligned} \hat{T}(\omega, r, m_q) |h, n, \mathbf{k}\rangle_{(r, m_q)}^{(\omega)} &= \\ e^{-a E_{h, n}(\mathbf{k}; \omega, r, m_q)} |h, n, \mathbf{k}\rangle_{(r, m_q)}^{(\omega)}. \end{aligned} \quad (16)$$

In the notations of ref. [1] one gets the formulae

$$\begin{aligned} E_{h, n}(\mathbf{k}; \omega, r, m_q) + E_{h, n}(\mathbf{k}; \omega, -r, m_q) &= \\ = 2E_{h, n}^{\text{cont}}(\mathbf{k}; m_q) + \mathcal{O}(a^2), \end{aligned} \quad (17)$$

$$\begin{aligned} \left[\langle h, n, \mathbf{k} | B | h', n', \mathbf{k}' \rangle \Big|_{(r, m_q)}^{(\omega)} + (r \rightarrow -r) \right] &= \\ = 2\zeta_B^B(\omega, r) \langle h, n, \mathbf{k} | B | h', n', \mathbf{k}' \rangle \Big|_{(m_q)}^{\text{cont}} + \mathcal{O}(a^2), \end{aligned} \quad (18)$$

where B is a gauge invariant m.r. local operator. It is important to remark that in the whole argument about $\mathcal{O}(a)$ improvement of WA 's the twisting angle ω is a totally inert label.

The interesting observation is that at $\omega = \pi/2$ the second term of the WA 's in eqs. (17) and (18), i.e. the quantities evaluated with Wilson parameter $-r$, can be rewritten in terms of closely related

⁴Actually the transfer matrix has been constructed only for $|r| = 1$. For $|r| < 1$ one should make reference to its square [9]. For simplicity we do not do that in the following.

quantities but evaluated with the value r of the Wilson parameter. As a consequence, $\mathcal{O}(a)$ improved estimates of energies and matrix elements can be obtained without having to average results from simulations with lattice actions differing by the sign of the Wilson term and critical mass.

Before coming to the proof of this statement, we recall from ref. [1] that at generic values of ω the product

$$\mathcal{P} \times (\omega \rightarrow -\omega), \quad (19)$$

where $\mathcal{P}$ is the physical parity operator ($x_P = (-\mathbf{x}, t)$)

$$\mathcal{P} : \begin{cases} U_0(x) \rightarrow U_0(x_P), \\ U_k(x) \rightarrow U_k^\dagger(x_P - a\hat{k}), k = 1, 2, 3 \\ \psi(x) \rightarrow \gamma_0 \psi(x_P) \\ \bar{\psi}(x) \rightarrow \bar{\psi}(x_P) \gamma_0 \end{cases} \quad (20)$$

is a spurionic symmetry of the tm-LQCD action (1). Since $[\mathcal{P} \times (\omega \rightarrow -\omega)]^2 = \mathbb{1}$, (multi-local) m.r. operators can always be taken to have a definite parity, which can be read off from the formula

$$\langle O^{(p)}(\{x_i\}) \rangle \Big|_{(r, m_q)}^{(\omega)} = (-1)^p \langle O^{(p)}(\{x_{iP}\}) \rangle \Big|_{(r, m_q)}^{(-\omega)}.$$

This relation entails the possibility of defining a notion of parity for the eigenstates of the transfer matrix [1]. One can, in fact, prove the validity of the following equations

$$\hat{\mathcal{P}} \hat{T}(\omega, r, m_q) \hat{\mathcal{P}} = \hat{T}(-\omega, r, m_q), \quad (21)$$

$$\hat{\mathcal{P}} |h, n, \mathbf{k}\rangle_{(r, m_q)}^{(\omega)} = \eta_{h, n} |h, n, -\mathbf{k}\rangle_{(r, m_q)}^{(-\omega)}, \quad (22)$$

$$E_{h, n}(\mathbf{k}; \omega, r, m_q) = E_{h, n}(-\mathbf{k}; -\omega, r, m_q), \quad (23)$$

where $\hat{\mathcal{P}}$ is the representative of the parity operation on the Hilbert space of states of the theory and $\eta_{h, n}$ ($\eta_{h, n}^2 = 1$) is what we will call the parity of the state $|h, n, \mathbf{k}\rangle_{(r, m_q)}^{(\omega)}$. Indeed $\eta_{h, n}$ can be taken to be an ω -independent integer coinciding with the physical parity label of the corresponding continuum state. For details we refer the reader to Appendix F of ref. [1].

It is also immediate to recognize that the action (1) goes into itself under the transformation (we recall that ω is defined mod 2π)

$$(r \rightarrow -r) \times (\omega \rightarrow \omega \pm \pi). \quad (24)$$

If in particular we set, say, $\omega = \pi/2$, the tm-LQCD action takes the form (4) and the invariance (24) becomes

$$(r \rightarrow -r) \times (\omega \rightarrow -\omega)|_{\omega=\pi/2}. \quad (25)$$

Either by inspection or by observing that, owing to (25), (19) is equivalent to

$$\mathcal{P} \times (r \rightarrow -r), \quad (26)$$

we conclude that the action (4) is invariant under (26). This means that at $\omega = \pi/2$ eq. (16) and equations from (21) to (23) can be rewritten in the form (for short we drop the twisting angle label)

$$\left[\widehat{T}(r, m_q) - e^{-aE_{h,n}(\mathbf{k}; r, m_q)} \right] |h, n, \mathbf{k}\rangle \Big|_{(r, m_q)} = 0 \quad (27)$$

$$\widehat{\mathcal{P}} \widehat{T}(r, m_q) \widehat{\mathcal{P}} = \widehat{T}(-r, m_q), \quad (28)$$

$$\widehat{\mathcal{P}} |h, n, \mathbf{k}\rangle \Big|_{(r, m_q)} = \eta_{h,n} |h, n, -\mathbf{k}\rangle \Big|_{(-r, m_q)}, \quad (29)$$

$$E_{h,n}(\mathbf{k}; r, m_q) = E_{h,n}(-\mathbf{k}; -r, m_q). \quad (30)$$

Using the last two equations it is now possible to cast at $\omega = \pi/2$ the WA 's (18) and (17) in a form in which all the relevant lattice data are extracted from simulations carried out with the action (4) and a given fixed value, r , of the Wilson parameter. One gets, in fact, the formulae

$$\begin{aligned} & \langle h, n, \mathbf{k} | B | h', n', \mathbf{k}' \rangle \Big|_{(r, m_q)} + \\ & + \eta_{hnh'n'}^B \langle h, n, -\mathbf{k} | B | h', n', -\mathbf{k}' \rangle \Big|_{(r, m_q)} = \\ & = 2\zeta_B^B(r) \langle h, n, \mathbf{k} | B | h', n', \mathbf{k}' \rangle \Big|_{(m_q)}^{\text{cont}} + \mathcal{O}(a^2), \quad (31) \end{aligned}$$

where $\eta_{hnh'n'}^B = \eta_{hn}(-1)^{p_B} \eta_{h'n'} = \pm 1$, and

$$\begin{aligned} & E_{h,n}(\mathbf{k}; r, m_q) + E_{h,n}(-\mathbf{k}; r, m_q) = \\ & = 2E_{h,n}^{\text{cont}}(\mathbf{k}; m_q) + \mathcal{O}(a^2). \quad (32) \end{aligned}$$

We observe that $\eta_{hnh'n'}^B$ is the product of the parities of the states $|h, n, \mathbf{k}\rangle$ and $|h', n', \mathbf{k}'\rangle$ (which, if not known, can be determined numerically as explained in ref. [1]) times the parity, $(-1)^{p_B}$, of the local m.r. operator B .

Notice that in case the lattice matrix element $\langle h, n, \mathbf{k} | B | h', n', \mathbf{k}' \rangle|_{(r, m_q)}$ is invariant under inversion of all the external three-momenta (like when all three-momenta vanish), the formula (31) gets particularly simple. In fact, if $\eta_{hnh'n'}^B = 1$, the lattice matrix element turns out to be automatically $\mathcal{O}(a)$ improved, while if $\eta_{hnh'n'}^B = -1$ eq. (31) implies that the $\mathcal{O}(a)$ improved estimate of $\langle h, n, \mathbf{k} | B | h', n', \mathbf{k}' \rangle|_{(r, m_q)}$ is zero. This last result is in agreement with what one expects in the continuum limit from parity invariance.

An important instance of a quantity which is automatically $\mathcal{O}(a)$ improved is F_π [1], as it can be extracted from two-point correlators evaluated at zero external three-momentum.

Eq. (32) taken at $\mathbf{k} = \mathbf{0}$ tells us that masses are automatically $\mathcal{O}(a)$ improved.

We end by observing that, although hadronic masses are automatically $\mathcal{O}(a)$ improved at $\omega = \pm\pi/2$, it is not possible to extract M_{cr} from numerical simulations with a discretization error which is better than $\mathcal{O}(a)$ (unless the theory is fully $\mathcal{O}(a)$ improved à la Symanzik). The reason is precisely that at $\omega = \pm\pi/2$ the pion mass is automatically $\mathcal{O}(a)$ improved, even if the critical mass is known with an $\mathcal{O}(a)$ error (see Appendix D of ref. [1]), thus it cannot be sensitive to $\mathcal{O}(a)$ discretization uncertainties in the critical mass. With an analogous argument a similar conclusion is reached if WTI's are used to determine M_{cr} .

3. Mass non-degenerate quarks

The maximally twisted LQCD fermionic action of an $SU_f(2)$ pair of mass non-degenerate quark can be conveniently written in the form

$$\begin{aligned} S_{\text{F,ND}}^{(\pi/2)}[\psi, \bar{\psi}, U] &= a^4 \sum_x \bar{\psi}(x) \left[\gamma \cdot \tilde{\nabla} + \right. \quad (33) \\ & \left. - i\gamma_5 \tau_1 W_{\text{cr}}(r) + m_q + \tau_3 \epsilon_q \right] \psi(x), \end{aligned}$$

where to keep the mass term real and flavour diagonal we have used the matrix τ_3 to split the masses of the members of the doublet. Consequently the Wilson term was twisted with the flavour matrix τ_1 . Our notations are such that m_q and ϵ_q are both positive.

What we said about the renormalizability of the fermion action in the mass degenerate case is valid also here and it is explicitly proved in Appendix A.

3.1. Non-singlet Ward-Takahashi identities

For mass non-degenerate quarks non-singlet WTI's take the continuum-like form ($x \neq y$)

$$\begin{aligned} \langle [\partial_\mu^* \hat{V}_\mu^1(x) - 2i\hat{\epsilon}_q \hat{S}^2(x)] \hat{O}(y) \rangle \Big|_{(r, m_q, \epsilon_q)} &= O(a) \\ \langle [\partial_\mu^* \hat{V}_\mu^2(x) + 2i\hat{\epsilon}_q \hat{S}^1(x)] \hat{O}(y) \rangle \Big|_{(r, m_q, \epsilon_q)} &= O(a) \\ \langle \partial_\mu^* \hat{V}_\mu^3(x) \hat{O}(y) \rangle \Big|_{(r, m_q, \epsilon_q)} &= O(a) \\ \langle [\partial_\mu^* \hat{A}_\mu^1(x) - 2\hat{m}_q \hat{P}^1(x)] \hat{O}(y) \rangle \Big|_{(r, m_q, \epsilon_q)} &+ O(a) \\ \langle [\partial_\mu^* \hat{A}_\mu^2(x) - 2\hat{m}_q \hat{P}^2(x)] \hat{O}(y) \rangle \Big|_{(r, m_q, \epsilon_q)} &= O(a) \\ \langle [\partial_\mu^* \hat{A}_\mu^3(x) - 2\hat{m}_q \hat{P}^3(x) + \\ -\hat{\epsilon}_q \hat{P}^0(x)] \hat{O}(y) \rangle \Big|_{(r, m_q, \epsilon_q)} &= O(a), \end{aligned}$$

if we make use of the definitions

$$\begin{aligned} \hat{V}_\mu^1 &= Z_V \bar{\psi} \gamma_\mu \frac{\tau_1}{2} \psi & \hat{A}_\mu^1 &= Z_A \bar{\psi} \gamma_\mu \gamma_5 \frac{\tau_1}{2} \psi \\ \hat{V}_\mu^2 &= Z_A \bar{\psi} \gamma_\mu \frac{\tau_2}{2} \psi & \hat{A}_\mu^2 &= Z_V \bar{\psi} \gamma_\mu \gamma_5 \frac{\tau_2}{2} \psi \\ \hat{V}_\mu^3 &= Z_A \bar{\psi} \gamma_\mu \frac{\tau_3}{2} \psi & \hat{A}_\mu^3 &= Z_V \bar{\psi} \gamma_\mu \gamma_5 \frac{\tau_3}{2} \psi \end{aligned} \quad (34)$$

$$\hat{P}^1 = Z_{S^0} \left[\bar{\psi} \frac{\tau_1}{2} \gamma_5 \psi + a^{-3} i \rho_P(am_q, a\epsilon_q) \mathbb{1} \right] \quad (35)$$

$$\hat{P}^b = Z_P \bar{\psi} \frac{\tau_b}{2} \gamma_5 \psi, \quad b = 2, 3 \quad (36)$$

$$\hat{P}^0 = Z_S \bar{\psi} \gamma_5 \psi \quad (37)$$

$$\hat{S}^1 = Z_{P^0} \bar{\psi} \frac{\tau_1}{2} \psi \quad (38)$$

$$\hat{S}^2 = Z_S \bar{\psi} \frac{\tau_2}{2} \psi \quad (39)$$

$$\hat{m}_q = Z_P^{-1} m_q \quad \hat{\epsilon}_q = Z_S^{-1} \epsilon_q. \quad (40)$$

As expected, the above formulae turn into eqs. (6) to (11) if we set $\epsilon_q = 0$ and perform the cyclic permutation of flavour indices $3 \rightarrow 2 \rightarrow 1 \rightarrow 3$. The above WTI's allow us to identify

$$\hat{m}_q^{(+)} = \hat{m}_q + \hat{\epsilon}_q = Z_P^{-1} m_q + Z_S^{-1} \epsilon_q, \quad (41)$$

$$\hat{m}_q^{(-)} = \hat{m}_q - \hat{\epsilon}_q = Z_P^{-1} m_q - Z_S^{-1} \epsilon_q \quad (42)$$

as the renormalized masses of the quarks in the doublet.

For completeness we record the formulae

$$\hat{S}^0 = Z_P \left[\bar{\psi} \psi + a^{-2} m_q \rho_{S^0}(am_q, a\epsilon_q) \mathbb{1} \right] \quad (43)$$

$$\hat{S}^3 = Z_S \left[\bar{\psi} \frac{\tau_3}{2} \psi + a^{-2} \epsilon_q \rho_S(am_q, a\epsilon_q) \mathbb{1} \right], \quad (44)$$

where $\rho_{S^0}(am_q, a\epsilon_q)$ and $\rho_S(am_q, a\epsilon_q)$, as well as $\rho_P(am_q, a\epsilon_q)$, are dimensionless real coefficients that admit a polynomial expansion both in am_q and $a\epsilon_q$. The mixing coefficients $\rho_P(am_q)$ and $\rho_{S^0}(am_q)$, appearing in eqs. (11) and (12) in the mass degenerate case, coincide with the value at $\epsilon_q = 0$ of the coefficients introduced in eq. (35) and (43), respectively.

3.2. O(a) improvement

The method for O(a) improvement is just as in the mass degenerate case. The physical explanation of this fact is obvious: all O(a) discretization effects come from the Wilson term which is of the same form in the actions (4) and (33) (up to a trivial flavour rotation). Formally, it is enough to observe that the action (33) is invariant under the transformations $\mathcal{R}_5 \times \mathcal{D}_d$ and

$$\begin{aligned} \mathcal{R}_5^{\text{spND}} &\equiv \\ &\equiv \mathcal{R}_5 \times (r \rightarrow -r) \times (m_q \rightarrow -m_q) \times (\epsilon_q \rightarrow -\epsilon_q), \end{aligned} \quad (45)$$

as well as under the spurionic parity operation (26). Invariance under $\mathcal{R}_5^{\text{spND}}$ and $\mathcal{R}_5 \times \mathcal{D}_d$ allows to prove (by arguments analogous to those presented in ref. [1]) that WA's of energies and matrix elements are free from O(a) cutoff effects. In formulae (dropping the label $\omega = \pi/2$)

$$\begin{aligned} E_{h,n}(\mathbf{k}; r, m_q, \epsilon_q) + E_{h,n}(\mathbf{k}; -r, m_q, \epsilon_q) &= \\ &= 2E_{h,n}^{\text{cont}}(\mathbf{k}; m_q, \epsilon_q) + O(a^2), \end{aligned} \quad (46)$$

$$\begin{aligned} \left[\langle h, n, \mathbf{k} | B | h', n', \mathbf{k}' \rangle \right]_{(r, m_q, \epsilon_q)} + (r \rightarrow -r) &= \\ &= 2\zeta_B^B(r) \langle h, n, \mathbf{k} | B | h', n', \mathbf{k}' \rangle \Big|_{(m_q, \epsilon_q)}^{\text{cont}} + O(a^2). \end{aligned} \quad (47)$$

Using the symmetry of the action (33) under the transformation (26), one can then prove, along the lines of sect. 2.2, that averages of ener-

gies evaluated with opposite values of the three-momenta as well as the appropriate linear combinations of matrix elements of m.r. operators taken between pairs of states with opposite external three-momenta are not affected by $O(a)$ discretization errors. In formulae

$$\begin{aligned} E_{h,n}(\mathbf{k}; r, m_q, \epsilon_q) + E_{h,n}(-\mathbf{k}; r, m_q, \epsilon_q) = \\ = 2E_{h,n}^{\text{cont}}(\mathbf{k}; m_q, \epsilon_q) + O(a^2), \end{aligned} \quad (48)$$

$$\begin{aligned} \langle h, n, \mathbf{k} | B | h', n', \mathbf{k}' \rangle \Big|_{(r, m_q, \epsilon_q)} + \\ + \eta_{hh'n'}^B \langle h, n, -\mathbf{k} | B | h', n', -\mathbf{k}' \rangle \Big|_{(r, m_q, \epsilon_q)} = \\ = 2\zeta_B^B(r) \langle h, n, \mathbf{k} | B | h', n', \mathbf{k}' \rangle \Big|_{(m_q, \epsilon_q)}^{\text{cont}} + O(a^2). \end{aligned} \quad (49)$$

3.3. The fermion determinant

The fermion determinant associated with the action (33) is real and strictly positive, provided

$$m_q^2 > \epsilon_q^2. \quad (50)$$

The proof, though elementary, requires some algebra and is presented in detail in Appendix B.

It is important to realize that (50) is not a trivial condition, because in terms of the renormalized quark masses, $\hat{m}_q^{(\pm)}$, it implies the inequality

$$\frac{Z_P}{Z_S} > \frac{\hat{m}_q^{(+)} - \hat{m}_q^{(-)}}{\hat{m}_q^{(+)} + \hat{m}_q^{(-)}}. \quad (51)$$

The latter leads to a rather stringent constraint on the (finite) ratio Z_P/Z_S , if $\hat{m}_q^{(+)} \gg \hat{m}_q^{(-)}$. One gets, in fact

$$\frac{Z_P}{Z_S} > 1 - 2 \frac{\hat{m}_q^{(-)}}{\hat{m}_q^{(+)}} + O\left[\left(\frac{\hat{m}_q^{(-)}}{\hat{m}_q^{(+)}}\right)^2\right]. \quad (52)$$

Since numerically one finds [10,11] for the quenched ratio Z_P/Z_S a number somewhat smaller than 1, one might be worried that the inequality (51) is not fulfilled. On this issue a leverage can be offered by the choice of the value of r which can be any real number satisfying $0 < |r| \leq 1$. In perturbation theory one finds for the standard Wilson action to one-loop [6] something like $Z_P/Z_S = 1 - g_0^2 r^2 I(r) + \dots$, where

$I(r) = I(-r) > 0$ for $0 < |r| \leq 1$ and has a finite limit at $r = 0$. This suggests that decreasing $|r|$ may increase Z_P/Z_S .

4. Conclusions

In this note we have shown that tm-LQCD at $\omega = \pm\pi/2$ yields a particularly useful lattice regularization of the Wilson type for the gauge theory of mass non-degenerate fermion pairs. Following ref. [1], we have proved that with this action essentially all physically relevant quantities can be evaluated with no $O(a)$ cutoff effects, while at the same time having the fermion determinant real and strictly positive, provided $0 < \epsilon_q^2 < m_q^2$. Monte Carlo simulations are hence safely feasible. The difficulties of the standard HMC in the presence of a flavour non-diagonal structure of the WD operator, D_{ND} (see eq. (67)), can be overcome e.g. by stochastically computing its determinant using algorithms of the multi-boson [12] or PHMC [13] type. Such algorithms should now be based on some polynomial approximation of $1/\sqrt{D_{\text{ND}}^\dagger D_{\text{ND}}}$ and naturally allow a simple correction of the employed polynomial approximation.

The approach we have discussed can be adapted [5] to the calculation of matrix elements of the CP-conserving, $\Delta S = 2$ and $\Delta S = 1$ effective weak Hamiltonian, in a way which we expect will solve the problem of “wrong chirality mixings” in the construction of the corresponding m.r. lattice operators.

Acknowledgements - One of us (G.C.R.) would like to thank the organizers of LHP2003 for the lively and exciting atmosphere of the Conference and its remarkably good organization. This work was supported in part by the European Community’s Human Potential Programme under contract HPRN-CT-2000-00145, Hadrons/Lattice QCD.

Appendix A: Symmetries of the action (33) and absence of counter-terms

We want to show that the (spurionic) symmetries enjoyed by the fermionic action (33) ensure

that no extra operators (of dimension ≤ 4) can be generated by radiative corrections, besides those already present. The discussion that follows applies in particular to the case $\epsilon_q = 0$, implying the “stability” under radiative corrections of the action (4) ⁵.

The list of transformations which leave the action (33) invariant include

1) lattice gauge transformations, space-time translations and hyper-cubic rotations, as well as the $U_V(1)$ vector transformation associated to baryon number conservation (none of these will play any special role in the discussion)

2) three continuous (one vector, $I^1(\theta_1)$, and two axial, $I^2(\theta_2)$, $I^3(\theta_3)$) non-singlet spurionic transformations

$$\mathcal{I}^1(\theta_1) \times (m_q + \epsilon_q \tau_3 \rightarrow e^{i\theta_1 \tau_1/2} [m_q + \epsilon_q \tau_3] e^{-i\theta_1 \tau_1/2}), \quad (53)$$

$$\mathcal{I}^2(\theta_2) \times (m_q + \epsilon_q \tau_3 \rightarrow e^{i\theta_2 \gamma_5 \tau_3/2} [m_q + \epsilon_q \tau_3] e^{i\theta_2 \gamma_5 \tau_3/2}), \quad (54)$$

$$\mathcal{I}^3(\theta_3) \times (m_q + \epsilon_q \tau_3 \rightarrow e^{-i\theta_3 \gamma_5 \tau_2/2} [m_q + \epsilon_q \tau_3] e^{-i\theta_3 \gamma_5 \tau_2/2}), \quad (55)$$

where (notice the simplification occuring if $b = 1$)

$$\mathcal{I}^b(\theta) : \begin{cases} \psi(x) \rightarrow e^{i\gamma_5 \tau_1 \pi/4} e^{i\theta \tau_b/2} e^{-i\gamma_5 \tau_1 \pi/4} \psi(x) \\ \bar{\psi}(x) \rightarrow \bar{\psi}(x) e^{-i\gamma_5 \tau_1 \pi/4} e^{-i\theta \tau_b/2} e^{i\gamma_5 \tau_1 \pi/4} \end{cases}$$

3) charge conjugation, $\mathcal{C}$

$$\mathcal{C} : \begin{cases} U_\mu(x) \rightarrow U_\mu^\star(x) \\ \psi(x) \rightarrow i\gamma_0 \gamma_2 \bar{\psi}^T(x) \\ \bar{\psi}(x) \rightarrow -i\psi^T(x) \gamma_0 \gamma_2 \end{cases} \quad (56)$$

4) the (anti-linear) reflection operation, Θ . It can be defined either with respect to the time slice $x_0 = 0$ (site reflection, Θ_s) or $x_0 = a/2$ (link reflection, Θ_ℓ). Its action on monomials of fermionic fields is

$$\begin{aligned} \Theta[f(U)\psi(x_1)\dots\bar{\psi}(x_n)] &= \\ &= f^\star(\Theta[U])\Theta[\bar{\psi}(x_n)]\dots\Theta[\psi(x_1)], \end{aligned} \quad (57)$$

where $f(U)$ is a functional of link variables and

$$\Theta_{s/\ell}[U_k(x)] = U_k^\star(\theta_{s/\ell}x),$$

⁵The action (33) at $\epsilon_q = 0$ is related to the action (4) by the harmless flavour rotation $\psi \rightarrow \exp(-i\tau_2 \pi/4)\psi$, $\bar{\psi} \rightarrow \bar{\psi} \exp(i\tau_2 \pi/4)$.

$$\Theta_{s/\ell}[U_0(x)] = U_0^T(\theta_{s/\ell}x - a\hat{0}), \quad (58)$$

$$\Theta_{s/\ell}[\psi(x)] = \bar{\psi}(\theta_{s/\ell}x)\gamma_0,$$

$$\Theta_{s/\ell}[\bar{\psi}(x)] = \gamma_0\psi(\theta_{s/\ell}x), \quad (59)$$

with

$$\theta_\ell(\mathbf{x}, t) = (\mathbf{x}, -t + a), \quad \theta_s(\mathbf{x}, t) = (\mathbf{x}, -t) \quad (60)$$

5) the pseudo-parity transformations

$$\mathcal{P}_{\pi/2}^1 \times (m_q \rightarrow -m_q), \mathcal{P}_F^2 \times (\epsilon_q \rightarrow -\epsilon_q), \mathcal{P}_F^3, \quad (61)$$

where ($x_P \equiv (-\mathbf{x}, x_0)$)

$$\mathcal{P}_{\pi/2}^1 : \begin{cases} U_0(x) \rightarrow U_0(x_P), \\ U_k(x) \rightarrow U_k^\dagger(x_P - a\hat{k}), k = 1, 2, 3 \\ \psi(x) \rightarrow i\gamma_5 \tau_1 \gamma_0 \psi(x_P) \\ \bar{\psi}(x) \rightarrow i\bar{\psi}(x_P) \gamma_0 \gamma_5 \tau_1 \end{cases} \quad (62)$$

$$\mathcal{P}_F^{2,3} : \begin{cases} U_0(x) \rightarrow U_0(x_P), \\ U_k(x) \rightarrow U_k^\dagger(x_P - a\hat{k}), k = 1, 2, 3 \\ \psi(x) \rightarrow i\tau_{2,3} \gamma_0 \psi(x_P) \\ \bar{\psi}(x) \rightarrow -i\bar{\psi}(x_P) \gamma_0 \tau_{2,3} \end{cases} \quad (63)$$

6) the transformation $\mathcal{R}_5 \times \mathcal{D}_d$, where

$$\mathcal{R}_5 : \begin{cases} \psi(x) \rightarrow \gamma_5 \psi(x) \\ \bar{\psi}(x) \rightarrow -\bar{\psi}(x) \gamma_5 \end{cases} \quad (64)$$

$$\mathcal{D}_d : \begin{cases} U_\mu(x) \rightarrow U_\mu^\dagger(-x - a\hat{\mu}) \\ \psi(x) \rightarrow e^{3i\pi/2} \psi(-x) \\ \bar{\psi}(x) \rightarrow e^{3i\pi/2} \bar{\psi}(-x) \end{cases} \quad (65)$$

7) the spurionic transformations (26), (45) and $\mathcal{T}_E \times (r \rightarrow -r)$, where $\mathcal{T}_E$ is the (euclidean) time-axis inversion. The action of $\mathcal{T}_E$ on the elementary fields of the theory is ($x_T \equiv (\mathbf{x}, -x_0)$)

$$\mathcal{T}_E : \begin{cases} U_0(x) \rightarrow U_0^\dagger(x_T - a\hat{0}), \\ U_k(x) \rightarrow U_k(x_T), k = 1, 2, 3 \\ \psi(x) \rightarrow \gamma_5 \gamma_0 \psi(x_T) \\ \bar{\psi}(x) \rightarrow \bar{\psi}(x_T) \gamma_0 \gamma_5 \end{cases} \quad (66)$$

In Tables 1 and 2 we list all the independent operators of dimension not larger than 4 that cannot appear in the action (33). They are grouped in columns under the name of the corresponding “killing” symmetry. The reality properties of the various coefficients in the action (33) are fixed by the invariance under the anti-linear reflection operation Θ_ℓ (or Θ_s).

Table 1

We list under the name of the relevant symmetry that forbids them all the operators of dimension $d = 4$ and $d = 3$ that cannot appear in the density action (33). By $F\tilde{F}$ we mean any lattice discretization of $\epsilon_{\mu\nu\lambda\rho}\text{Tr}[F_{\mu\nu}F_{\lambda\rho}]$.

Dim	$\mathcal{C}$
$d = 4$	$\bar{\psi}\mathcal{T}\gamma_5\gamma\tilde{\nabla}\psi$ ($\mathcal{T} = 1, \tau_1, \tau_3$)
$d = 3$	$\bar{\psi}\tau_2\psi$ $\bar{\psi}\tau_2\gamma_5\psi$

Table 2

Continuation of Table 1

Dim	P_F^3	$P_F^2 \times (\epsilon_q \rightarrow -\epsilon_q)$
$d = 4$	$\bar{\psi}\mathcal{T}\gamma\tilde{\nabla}\psi$ ($\mathcal{T} = \tau_1, \tau_2$) $F\tilde{F}$	$\bar{\psi}\tau_2\gamma_5\gamma\tilde{\nabla}\psi$ $\bar{\psi}\tau_3\gamma\tilde{\nabla}\psi$
$d = 3$	$\bar{\psi}\gamma_5\psi$ $\bar{\psi}\gamma_5\tau_3\psi$ $\bar{\psi}\tau_1\psi$	

The conclusion of this discussion is that the form (33) of the action is preserved by radiative corrections and the bare parameters, g_0^2 , m_q and ϵ_q , need only a purely multiplicative renormalization. More details on the renormalization of m_q and ϵ_q can be found in the text in sect. 3.1.

Appendix B: Positivity of the fermion determinant

In this Appendix we show that under the condition (50) the determinant of the WD operator associated with the action (33), namely

$$D_{\text{ND}} = \gamma \cdot \tilde{\nabla} - i\gamma_5\tau_1 W_{\text{cr}}(r) + m_q + \tau_3\epsilon_q, \quad (67)$$

is real and (strictly) positive. To prove that $D_{\text{F,ND}} = \text{Det}[D_{\text{ND}}]$ is a real number it is enough

to note the self-adjointness relation

$$\gamma_5\tau_3 D_{\text{ND}}\gamma_5\tau_3 = D_{\text{ND}}^\dagger. \quad (68)$$

The proof of positivity is somewhat more involved. To proceed it is convenient to introduce the auxiliary self-adjoint operator

$$Q_{\text{cr}} = \gamma_5 \left[\gamma \cdot \tilde{\nabla} + W_{\text{cr}}(r) \right] = Q_{\text{cr}}^\dagger, \quad (69)$$

in terms of which $\mathcal{D}_{\text{F,ND}}$ can be written in the form

$$\mathcal{D}_{\text{F,ND}} = \text{Det}[Q_{\text{cr}} + i\tau_1 m_q + \gamma_5\tau_3\epsilon_q]. \quad (70)$$

In flavour space the operator in the r.h.s. of eq. (70) is represented by the 2×2 matrix

$$Q_{\text{cr}} + i\tau_1 m_q + \gamma_5\tau_3\epsilon_q = \gamma_5 e^{i\pi\gamma_5\tau_1/4} D_{\text{ND}} e^{i\pi\gamma_5\tau_1/4} \\ = \begin{pmatrix} Q_{\text{cr}} + \gamma_5\epsilon_q & im_q \\ im_q & Q_{\text{cr}} - \gamma_5\epsilon_q \end{pmatrix}. \quad (71)$$

For the determinant of this operator one finds

$$\begin{aligned} \mathcal{D}_{\text{F,ND}} &= \text{Det}[D_{\text{ND}}] = \\ &= \det[(Q_{\text{cr}} + \gamma_5\epsilon_q)(Q_{\text{cr}} - \gamma_5\epsilon_q) + m_q^2] = \\ &= \det[Q_{\text{cr}}^2 + m_q^2 - \epsilon_q^2 + 2\epsilon_q\gamma \cdot \tilde{\nabla}] = \\ &= \det[Q_{\text{cr}}^2 + m_q^2 - \epsilon_q^2] \cdot \det[1 + 2\epsilon_q B], \end{aligned} \quad (72)$$

where we have introduced the definition

$$B = X \gamma \cdot \tilde{\nabla} X, \quad X = (Q_{\text{cr}}^2 + m_q^2 - \epsilon_q^2)^{-1/2}. \quad (73)$$

Since $\det[Q_{\text{cr}}^2 + m_q^2 - \epsilon_q^2] > 0$, we only have to prove that the second factor in the last equality of (72) is a positive quantity. To this end we observe that both $\mathcal{D}_{\text{F,ND}}$ (from eq. (71)) and $\det[Q_{\text{cr}}^2 + m_q^2 - \epsilon_q^2]$ (by inspection) are even under $\epsilon_q \rightarrow -\epsilon_q$. We then conclude that

$$\Delta_{\text{F,ND}}(\epsilon_q) \equiv \det[1 + 2\epsilon_q B] = \Delta_{\text{F,ND}}(-\epsilon_q). \quad (74)$$

This property is indeed enough to prove the positivity of $\Delta_{\text{F,ND}}$ and hence our thesis. Let us consider, in fact, the expansion

$$\text{tr}[\log(1 + 2\epsilon_q B)] = \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} (2\epsilon_q)^k \text{tr}[B^k], \quad (75)$$

valid for sufficiently small values of ϵ_q ⁶. Eq. (74) implies that only even powers of ϵ_q can contribute. This fact together with the observation

⁶For instance, one can take $2|\epsilon_q|$ smaller than the number $\min[m_q, 1/||B||_{2\epsilon_q=m_q}]$.

that B is an anti-Hermitian operator ($B^\dagger = -B$, because $\gamma \cdot \tilde{\nabla}$ is anti-Hermitian) allows us to write

$$\begin{aligned} \text{l.h.s. of (75)} &= \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{2n} (2\epsilon_q)^{2n} \text{tr}[(B^\dagger B)^n] = \\ &= \frac{1}{2} \text{tr}[\log(1 + 4\epsilon_q^2 B^\dagger B)]. \end{aligned} \quad (76)$$

The last equality proves that $\text{tr}[\log(1 + 2\epsilon_q B)]$ is a non-negative quantity, implying that $\det[1 + 2\epsilon_q B]$ is real and strictly positive.

We conclude with two technical observations:

1) since B is anti-Hermitian, its spectrum is purely imaginary, thus $1 + 2\epsilon_q B$ cannot have any vanishing eigenvalue; 2) eq. (76), which was proved for sufficiently small ϵ_q , can be extended to the actual physical value of the mass splitting by analyticity (see the chain of equalities in eq. (72)).

REFERENCES

1. R. Frezzotti and G.C. Rossi, hep-lat/0306014 and hep-lat/0309157.
2. W.A. Bardeen, A. Duncan, E. Eichten, G. Hockney and H. Thacker, Phys. Rev. **D57** (1998) 1633;
W.A. Bardeen, A. Duncan, E. Eichten and H. Thacker, Phys. Rev. **D59** (1999) 014507;
G. Schierholz et al., Nucl. Phys. **B** Proc. Suppl. **73** (1999) 889.
3. A.C. Irving, C. McNeile, C. Michael, K.J. Sharkey and H. Wittig [UKQCD Collaboration], Phys. Lett. **518B** (2001) 243.
4. R. Frezzotti, P.A. Grassi, S. Sint and P. Weisz, Nucl. Phys. **B** (Proc. Suppl.) **83** (2000) 941 and JHEP **0108** (2001) 058;
R. Frezzotti, Nucl. Phys. **B** (Proc. Suppl.) **119** (2003) 140, hep-lat/0210007;
R. Frezzotti, S. Sint and P. Weisz, JHEP **0107** (2001) 048;
M. Della Morte, R. Frezzotti, J. Heitger and S. Sint [ALPHA collaboration], JHEP **0110** (2001) 041;
R. Frezzotti and S. Sint, Nucl. Phys. **B** (Proc. Suppl.) **106** (2002) 814, hep-lat/0110140.
5. R. Frezzotti and G.C. Rossi, in preparation.
6. M. Bochicchio, L. Maiani, G. Martinelli, G.C. Rossi and M. Testa, Nucl. Phys. **B262** (1985) 331.
7. M. Guagnelli, J. Heitger, C. Pena, S. Sint, and A. Vladikas, Nucl. Phys. **B** (Proc. Suppl.) **106** (2002) 320;
C. Pena, S. Sint, and A. Vladikas, Nucl. Phys. **B** (Proc. Suppl.) **119** (2003) 368, hep-lat/0209045.
8. M. Testa, JHEP **9804** (1998) 002.
9. M. Lüscher, Comm. Math. Phys. **54** (1977) 283;
K. Osterwalder and E. Seiler, Ann. of Phys. **110** (1978) 440;
G. Immirzi and K. Yoshida, Nucl. Phys. **B210** (1982) 499;
P. Menotti and A. Pelissetto, Com. Math. Phys. **113** (1987) 369 and SNS-15/87 preprint, Proc. of the Symposium on “*Field Theory on the Lattice*”, Seillac, France (1987).
10. T. Bhattacharya, R. Gupta, W.J. Lee and S.R. Sharpe, Phys. Rev. **D63** (2001) 074505, hep-lat/0009038.
11. V. Gimenez, L. Giusti, F. Rapuano and M. Talevi, Nucl. Phys. **B531** (1998) 429, hep-lat/9806006.
12. M. Lüscher, Nucl. Phys. **B418** (1994) 637, hep-lat/9311007.
13. P. de Forcrand and T. Takaishi, Nucl. Phys. **B** (Proc. Suppl.) **53** (1997) 968, hep-lat/9608093;
R. Frezzotti and K. Jansen, Phys. Lett. **402B** (1997) 328, hep-lat/9702016.